# Repeated exposure to low levels of the chemical warfare agent VX activates cell survival related gene in mouse brain

CPT Jose M. Pizarro, Ph.D., M.S.

Division of Neuroscience WRAIR

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to completing and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar DMB control number.	ion of information. Send comments arters Services, Directorate for Infor	regarding this burden estimate of mation Operations and Reports	or any other aspect of th , 1215 Jefferson Davis I	is collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE 01 OCT 2005		2. REPORT TYPE N/A		3. DATES COVERED		
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER				
Repeated exposure to low levels of the chemical warfare agent VX activates cell survival related gene in mouse brain mouse brain				5b. GRANT NUMBER		
activates cen sui vivai relateu gene in mouse brain mouse brain				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Division of Neuroscience Walter Reed Army Institute of Research				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release, distribution unlimited						
13. SUPPLEMENTARY NOTES  See also ADM001851, Proceedings of the 2003 Joint Service Scientific Conference on Chemical & Biological Defense Research, 17-20 November 2003., The original document contains color images.						
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC	17. LIMITATION OF	18. NUMBER	19a. NAME OF			
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	- ABSTRACT <b>UU</b>	OF PAGES  19	RESPONSIBLE PERSON	

**Report Documentation Page** 


Form Approved OMB No. 0704-0188

#### Introduction

- VX is an excitotoxic compound.
- Neurons may respond to toxicity by expressing genes associated with cell survival.
- Neurotrophins:
  - neuronal development
  - neuronal plasticity and remodeling
  - cell survival
- Brain derived neurotrophic factor (BDNF) is modulated:
  - traumatic brain injury
  - ischemia
  - toxin exposure
- BDNF may be important for neuronal plasticity, cell survival and remodeling following brain trauma.


#### Purpose


 Evaluate whether repeated exposure to low levels of VX affects genes associated with cell survival and neuronal plasticity.




### Animal Exposure

- Repeated low-level exposure
  - Three groups:
    - Saline controls
    - 0.2 LD<sub>50</sub>
    - 0.4 LD<sub>50</sub>
  - 1/day for 5 days/wk for 2 wk
- The LD<sub>50</sub> value for VX given subcutaneously (s.c.) in mice was 21 µg/kg (Boskovic, 1979).
- The nerve agent VX was dissolved in saline and injected s.c. in volume of 1 ml/kg.








#### In situ Hybridization


- RNA probes for BDNF were generated by in vitro transcription using Strip-EZ<sup>TM</sup> RNA.
- For antisense <sup>33</sup>P-UTP labeled BDNF RNA probes, a plasmid containing a 460 bp BDNF
  - -linearized with EcoRI
  - -transcribed with T7 polymerase
- For sense <sup>33</sup>P-UTP labeled BDNF RNA probes
  - -linearized with SalI
  - -transcribed with SP6 polymerase

#### In situ Hybridization (cont.)


- Coronal sections (14 μm) were hybridized overnight with <sup>33</sup>P-UTP labeled cRNA probe.
- The sections were then exposed to a series of high temperature washes in various dilutions of SSC.
- The sections were then air-dried and autoradiographed on Kodak BioMax MS.
- The film was analyzed using the Model GS-800 Calibrated Imaging Densitometer and the Quantity One software.




#### In situ hybridization cRNA controls




#### Spatio-Temporal Distribution of BDNF mRNA




### BDNF mRNA expression in the hippocampal region following repeated exposure to low levels of VX




## BDNF mRNA expression in the thalamic region following repeated exposure to low levels of VX



### BDNF mRNA expression in the cortical region following repeated exposure to low levels of VX



# BDNF mRNA expression in the hypothalamic and amygdaloid regions following repeated exposure to low levels of VX



#### Summary

- We examined the temporal profile of BDNF mRNA expression at 2 hr and 72 hr after repeated injections of either saline, 0.2LD<sub>50</sub> VX, or 0.4LD<sub>50</sub> VX using *in situ* hybridization analysis.
- Animals that were injected with 0.4 LD<sub>50</sub> VX showed elevated levels of BDNF mRNA expression 2 hr and 72 hr later, in areas of the hippocampus, lateral amygdala, cortex, hypothalamus and thalamus.
- In some brain regions, 0.2 LD<sub>50</sub> VX also increased BDNF expression at 2 hr, but not at 72 hr, postexposure.

### Acknowledgment

- WRAIR
  - Lumley LA
  - Saviolakis GA
  - Bah MJ
  - Alagappan A
  - Chang WE
  - Shah JD
  - Robison CL
  - Meyerhoff JL
  - Dawood MY
  - Karp A

- USAMRICD
  - CPT Midboe
  - Cerasoli D
  - Clark C
  - SPC D'Ambrozio

This research was funded by Task Area N (Low Dose CWNA)